

# THE GALACTIC ENVIRONMENT OF THE NEVIII ABSORBER TOWARD HE0226–4110

JOHN S. MULCHAEY

The Observatories of the Carnegie Institution of Washington, 813 Santa Barbara St., Pasadena, CA 91101, U.S.A.

AND

HSIAO-WEN CHEN

Department of Astronomy and Astrophysics and Kavli Institute for Cosmological Physics, University of Chicago, Chicago, IL, 60637, U.S.A.

Accepted for Publication in *ApJ Letters*

## ABSTRACT

We report the discovery of a small galaxy system in the vicinity of the NeVIII absorber at  $z = 0.20701$  toward HE0226–4110. The galaxy system consists of two  $0.25 L_*$  disk galaxies and a  $0.05 L_*$  galaxy all within  $\Delta v < 300 \text{ km s}^{-1}$  and  $\rho \leq 200 h^{-1}$  physical kpc of the absorber. We consider various scenarios for the origin of the NeVIII absorption, including photo-ionized gas from a AGN, a starburst driven wind, a hot intragroup medium, hot gas in a galaxy halo, and a conductive front produced by cool clouds moving at high speed through a hot medium. We argue that the conductive front scenario is most likely responsible for producing the NeVIII feature, because it is consistent with the observed galactic environment around the absorber and because it naturally explains the multi-phase nature of the gas and the kinematic signatures of the absorption profiles. Although our preferred scenario suggests that NeVIII may not be directly probing the warm-hot intergalactic medium, it does imply the existence of an extended hot confining medium around a disk galaxy that may contain a significant reservoir of baryons in the form of hot gas.

*Subject headings:* cosmology: observations—intergalactic medium—quasars: absorption lines—galaxies: halos

## 1. INTRODUCTION

A census of the observed baryons in the local universe falls far short of the amount required by Big Bang Nucleosynthesis and the cosmic microwave background (Fukugita et al. 1998). Cosmological simulations suggest most of the missing baryons are in the form of  $10^5$ – $10^7 \text{ K}$  gas that resides in low-density regions such as groups and filaments (Cen & Ostriker 1999; Davé et al. 2001). Considerable effort has been made in recent years to locate this “Warm-Hot Intergalactic Medium” (WHIM) (Bregman 2007), but such gas has largely eluded detection. Ultraviolet absorption lines in the spectra of background quasars are considered a sensitive probe of the WHIM (Verner et al. 1994; Mulchaey et al. 1996). The most commonly used feature is the OVI  $\lambda\lambda 1031.926, 1037.617 \text{ \AA}$  absorption doublet. While many OVI systems have now been identified (Tripp et al. 2000, 2008; Savage et al. 2002; Danforth & Shull 2005; Thom & Chen 2008a,b), recent observations (Tripp et al. 2008; Thom & Chen 2008b) suggest that  $\sim 30$ – $50\%$  of the observed OVI absorbers may result from photo-ionized gas and therefore may not be probing the WHIM.

The presence of other high ionization lines may help distinguish between photo-ionized and collisionally-ionized systems. In particular, the ultraviolet lines of NeVIII  $\lambda\lambda 770.5, 780.3$  are potentially powerful tracers of hot collisionally ionized gas because of the high ionization potential required to produce the ions (207.28 eV). However, only one intergalactic NeVIII absorber has been reported to date (Savage et al. 2005; Lehner et al. 2006). The observed NeVIII absorption is accompanied by lower ionization features. A detailed ionization analysis shows that the NeVIII is best explained by collisionally ionized warm gas, while the low ionization features most likely result from photo-ionization (Sav-

age et al. 2005). The relationship between the collisionally-ionized and photo-ionized components is unclear. The galactic environment of the NeVIII absorber is also unknown. Here, we report the results of a galaxy survey in the vicinity of the NeVIII absorber that allows us to better understand the nature of this system and to examine the effectiveness of NeVIII as a probe of the WHIM. We adopt a  $\Lambda$ CDM cosmology,  $\Omega_M = 0.3$  and  $\Omega_\Lambda = 0.7$ , with a dimensionless Hubble constant  $h = H_0/(100 \text{ Mpc s}^{-1} \text{ km}^{-1})$  throughout this Letter.

## 2. THE NEVIII ABSORBER AT $Z = 0.20701$ TOWARD HE0226–4110

Using high-resolution FUSE and STIS observations with respective spectral resolutions of  $\text{FWHM} \approx 25 \text{ km s}^{-1}$  and  $7 \text{ km s}^{-1}$ , Savage et al. (2005) reported the first detection of intergalactic NeVIII  $\lambda\lambda 770, 780$  absorption at  $z = 0.20701$  toward HE0226–4110, together with detections of absorption transitions due to HI, CIII, OIII, OIV, OVI, NIII, SiIII, SVI and possibly SV. Different ionization species exhibit different kinematic signatures in the absorption-line profiles, suggesting a multi-phase medium nature of the underlying absorbing gas. Specifically, the absorption transitions due to low-ionization species such as HI, CIII, and OIII are resolved into two relatively narrow components (Doppler parameter  $b \approx 15 \text{ km s}^{-1}$ ) separated by  $\Delta v \approx 30 \text{ km s}^{-1}$  (see also Thom & Chen 2008b). The observed HI column densities of the absorbing gas are  $\log N(\text{HI}) = 15.06 \pm 0.04$  for the blueshifted component and  $\log N(\text{HI}) = 14.89 \pm 0.05$  for the redshifted component. The OVI absorption doublet shows a broad component of  $b_{\text{OVI}} \approx 31 \pm 2 \text{ km s}^{-1}$  that centers roughly between the two components found in the low-ionization lines with  $\log N(\text{OVI}) = 14.37 \pm 0.03$ , while the NeVIII (and SVI) absorption lines appear to be blueshifted by  $\approx 7$ – $18 \text{ km s}^{-1}$  with respect to the OVI absorption doublet with  $\log N(\text{NeVIII}) = 13.9 \pm 0.1$  ( $\log N(\text{SVI}) = 12.8 \pm 0.1$ ) and  $b_{\text{NeVIII}} = 23 \pm 15$

Electronic address: mulchaey@ociw.edu  
Electronic address: hchen@oddjob.uchicago.edu

$\text{km s}^{-1}$  ( $b_{\text{SVI}} \approx 39 \pm 20 \text{ km s}^{-1}$ ).

Savage et al. (2005) conducted an extensive ionization analysis of the absorbing gas. Using the observed absorption strengths of CIII, OIII, NIII, and SiIII and assuming a mean background radiation field of  $J_{912} = 1.9 \times 10^{-23} \text{ ergs cm}^{-2} \text{ s}^{-1} \text{ Hz}^{-1} \text{ sr}^{-1}$  at  $z = 0.2$  from Haardt & Madau (1996), the authors found that the relative abundances of these low-ionization species are characterized by a photo-ionized cloud of metallicity  $\approx 1/3$  solar, temperature  $T \sim 2 \times 10^4 \text{ K}$ , density  $n_{\text{H}} = 2.6 \times 10^{-5} \text{ cm}^{-3}$ , neutral fraction  $f_{\text{H}^0} = 2.5 \times 10^{-4}$ , and size  $\approx 57 \text{ kpc}$ . But applying a photo-ionization model to explain the observed large column densities of OVI and NeVIII with the same assumed radiation field would require the gas density to be still lower,  $n_{\text{H}} = 4.5 \times 10^{-7} \text{ cm}^{-3}$  and the inferred total pathlength near  $\sim 11 \text{ Mpc}$ . Such a scenario was ruled out by Savage et al. (2005), because the Hubble flow broadening of the absorption over such a large path length would be much larger than that found in the observed line widths.

Savage et al. (2005) further considered a collisional ionization scenario, adopting the  $1/3$  solar metallicity inferred for the low-ionization species under the photo-ionization model. The authors derived a gas temperature of  $T = 5.4 \times 10^5 \text{ K}$ , assuming the gas is under collisional ionization equilibrium. At this temperature, the observed line widths of NeVIII and OVI would imply a turbulent velocity field of  $b_{\text{turbulent}} \sim 20 \text{ km s}^{-1}$ . At  $T \lesssim 6 \times 10^5 \text{ K}$ , however, cooling efficiency approaches a maximum. The gas is therefore expected to be in non-equilibrium conditions, undergoing rapid cooling at a somewhat lower temperature (e.g. Gnat & Sternberg 2007).

In summary, Savage et al. (2005) concluded that the  $z = 0.20701$  absorber toward HE0226–4110 is part of a multi-phase system. In their preferred model, the low ionization features originate in a photo-ionized component of  $T \sim 2 \times 10^4 \text{ K}$ , while the NeVIII and most of the OVI absorption originate in collisionally ionized gas of  $T \sim 5 \times 10^5 \text{ K}$ . However, the connection between the photo-ionized cool component and the collisionally ionized hot gas is unclear.

### 3. THE GALAXY SYSTEM AT $Z = 0.207$

To help determine the nature of the  $z = 0.20701$  absorption system, we have performed a spectroscopic survey of the galaxies in the field around HE0226–4110. Complete details of our observing program can be found in Chen & Mulchaey (2009). Here we provide a brief summary of the galaxy survey. Spectroscopic candidates were selected from optical images covering a  $\sim 28 \times 28 \text{ arcmin}^2$  region centered on the QSO. Candidate galaxies were observed at the Magellan telescopes using the IMACS and LDSS-3 spectrographs. The spectra were reduced using the COSMOS software package and redshifts measured by cross-correlating the extracted spectra and galaxy templates. The mean redshift measurement uncertainty is  $\Delta z \approx 0.0003$ . Our survey is nearly 100% complete for galaxies brighter than  $R = 23$  at angular distances  $\Delta\theta \leq 2'$  (corresponding to projected distances of  $\rho \leq 285 h^{-1}$  physical kpc at  $z = 0.20701$ ) and approximately 50% complete at  $R \leq 22$  out to  $\Delta\theta \leq 10'$  ( $\rho \leq 1.4 h^{-1}$  physical Mpc at  $z = 0.20701$ ).

Our spectroscopic survey revealed a small association of three galaxies in the vicinity of the NeVIII absorber. These galaxies are all within  $|\Delta v| < 300 \text{ km s}^{-1}$  and  $\rho \leq 200 h^{-1}$  physical kpc of the absorber. A summary of their properties is presented in Table 1. The optical spectra and an  $R$ -band image are presented in Figure 1.

Table 1 shows that the galaxy with the smallest projected distance from the QSO (A) is an underluminous disk galaxy of luminosity  $\sim 0.05 L_*$  (adopting  $M_R - 5 \log h = -20.4$  from Blanton et al. 2003). The projected distance between A and HE0226–4110 is  $\rho = 26.5 h^{-1}$  physical kpc. The two other objects are more luminous disk galaxies of  $\sim 0.25 L_*$ . These galaxies have impact parameters of  $\rho = 76.5 h^{-1}$  physical kpc (B) and  $\rho = 197.3 h^{-1}$  physical kpc (C), respectively. Galaxies A and B have emission-line spectra consistent with normal star-forming galaxies, while C exhibits absorption dominated spectral features that indicates a more evolved underlying stellar population.

We note that based on the completeness of our survey, we can rule out the presence of additional galaxies near the redshift of the absorber down to  $M_R - 5 \log h = -16.1$  (roughly  $0.02 L_*$ ) within  $\rho = 285 h^{-1}$  kpc of the QSO. Four additional galaxies of  $M_R - 5 \log h = -17.2$  to  $M_R - 5 \log h = -20.45$  are found within  $|\Delta v| < 300 \text{ km s}^{-1}$  of the absorber but these galaxies are located at  $\rho = 802 - 1500 h^{-1}$  physical kpc from the QSO line of sight (Chen & Mulchaey 2009). They are therefore unlikely to host the absorbing gas found in the QSO absorption-line data.

### 4. THE ORIGIN OF THE NEVIII ABSORBER

The combination of the results of our galaxy survey in the field around HE0226–4110 and the known properties of the absorbing gas from Savage et al. (2005) provides a unique opportunity to examine the relationship between the highly ionized warm gas and the galaxy environment. Below we consider whether the absorber is produced in individual galactic halos, intragroup medium in a galaxy group, or outflows from either a starburst galaxy or AGN. The close proximity of the NeVIII absorber to multiple galaxies is similar to what is found for some OVI absorbers (e.g. Stocke et al. 2006; Wakker & Savage 2009). It suggests that the absorbing gas is unlikely to originate in underdense filamentary regions, but is likely to be gravitationally bound to galaxies.

#### 4.1. Intragroup Medium

To assess whether the absorber is likely to be associated with an intragroup medium or a halo around a galaxy, we first estimate individual halo sizes of the galaxies based on their intrinsic luminosities. Previous halo occupation studies of field galaxies have shown a monotonic relationship between halo mass  $M_h$  and galaxy luminosity for galaxies fainter than  $L_*$  and located at the centers of their dark matter halos. Adopting the  $M_h$  vs.  $M_R$  relation of Tinker & Conroy (2009), we estimate that galaxies A, B, and C reside in halos of  $M_h = 10^{11}$ ,  $10^{11.5}$ , and  $10^{11.5} h^{-1} M_\odot$ , respectively. The corresponding halo radii  $R_h \equiv R_{200}^1$  are  $R_h = 93, 137$ , and  $137 h^{-1}$  kpc for A, B, C, respectively. The expected halo radii of these galaxies together with the projected separations between one another indicate that galaxies B and C do not share a common halo, whereas galaxies A and B (separated by  $\approx 50 h^{-1}$  projected physical kpc and  $\approx 300 \text{ km s}^{-1}$ ) are likely to share a common halo (based on the two-point correlation function, the probability of having two unrelated galaxies at this separation is  $\sim 0.5\%$ ; Zehavi et al. (2005)). Analogous to the Local Group, it is probably more appropriate to consider B and C as two separate halos, instead of a single virial “group”. With this

<sup>1</sup> We adopt  $R_{200}$  to represent the size of a dark matter halo, which corresponds to the radius within which the enclosed mean density is 200 times the background matter density.

TABLE 1  
SUMMARY OF GALAXIES ASSOCIATED WITH THE NeVIII ABSORBER AT  $z = 0.20701$  TOWARD HE0226–4110

| Galaxy ID | $\Delta\alpha$<br>(arcsec) | $\Delta\delta$<br>(arcsec) | $\Delta\theta$<br>(arcsec) | $\rho$<br>( $h^{-1}$ kpc) | $B$              | $R$              | $I$              | $z_{\text{spec}}$ | Spectral Type <sup>a</sup> | $M_R$<br>$-5 \log h$ |
|-----------|----------------------------|----------------------------|----------------------------|---------------------------|------------------|------------------|------------------|-------------------|----------------------------|----------------------|
| A         | −7.9                       | −7.9                       | 11.2                       | 26.5                      | $22.85 \pm 0.06$ | $21.94 \pm 0.04$ | $21.20 \pm 0.03$ | 0.2065            | 2                          | −17.2                |
| B         | −19.8                      | −25.3                      | 32.1                       | 76.5                      | $22.05 \pm 0.03$ | $20.29 \pm 0.01$ | $19.38 \pm 0.01$ | 0.2078            | 2                          | −18.9                |
| C         | −53.4                      | 63.5                       | 82.9                       | 197.3                     | $22.71 \pm 0.05$ | $20.53 \pm 0.01$ | $19.65 \pm 0.01$ | 0.2077            | 1                          | −18.9                |

<sup>a</sup> Spectral type of the galaxies: “1” indicates absorption-line dominated galaxies, and “2” indicates emission-line dominated galaxies. See also Figure 1.

interpretation of the mass distribution, it seems unlikely that the NeVIII absorber would be associated with a diffuse intra-group medium.

#### 4.2. Extended Hot Gaseous Halo

Given the projected distances from the QSO and the  $R_{200}$  estimates above, only A and B are viable hosts for the NeVIII absorber. Here, we consider the possibility that the absorber originates in an extended hot halo around one of these galaxies. Hot halos around individual galaxies are expected in the hierarchical galaxy formation paradigm. While such halos are commonly found around early-type galaxies (Fabbiano et al. 1992), the presence of extended hot halos around disk galaxies has not been confirmed (Rasmussen et al. 2009) with the exception of the Milky Way for which an extended hot halo is implied from several observations (Sembach et al. 2003; Fang et al. 2006; Bregman & Lloyd-Davies 2007).

We first consider the more massive galaxy B. The estimated halo mass of galaxy B would imply a virial temperature of  $T_{\text{vir}} \sim 4 \times 10^5$  K, similar to the gas temperature ( $T = 5.4 \times 10^5$  K) derived for the NeVIII absorber under the assumption of a collisional ionization equilibrium (Savage et al. 2005). It is therefore possible that the absorber originates in an extended hot halo around galaxy B.

If the NeVIII absorption is produced by an extended hot halo, we can use the observed properties of the absorption to estimate the mass of the hot gas. Savage et al. (2005) derive a total hydrogen column density of  $\log N(\text{H}) = 19.9$  for the absorber under the assumptions of collisional ionization equilibrium and metallicity  $[Z/\text{H}] = -0.5$ . Assuming a gaseous radius equal to  $R_{200}$ , the total path length is  $l \sim 227 h^{-1}$  kpc at the impact distance of  $\rho = 76.5 h^{-1}$  kpc. This path length combined with the inferred column density, suggests the total hydrogen density is  $n(\text{H}) \sim 1.1 \times 10^{-4} h \text{ cm}^{-3}$ . For a uniform density, the total mass in the hot gas within the virial radius is  $M_{\text{gas}} \sim 4 \times 10^{10} h^{-2} M_{\odot}$ . This mass is significantly greater than the mass in stars  $M_{*} \sim 4 \times 10^9 h^{-2} M_{\odot}$  as inferred by the broad-band spectral energy distribution of the galaxy. The difference would suggest that the hot gas revealed by the NeVIII absorption is the dominant baryonic component.

We have argued based on the mean space density of dwarf galaxies (see § 4.1) that galaxies A and B share a common halo. If instead A is at a cosmological distance from B and responsible for the absorption, the virial temperature is  $T_{\text{vir}} \sim 2 \times 10^5$  K for its host halo, somewhat lower than the temperature derived by Savage et al. (2005). The gas mass in this case would be  $M_{\text{gas}} \sim 9 \times 10^9 h^{-2} M_{\odot}$ . This mass is approximately a factor of 10 times larger than the stellar mass, so as is the case for galaxy B, the hot gas would be the dominant baryonic component if the NeVIII absorption was due to a hot gas halo.

However, the notion of NeVIII arising in an extended hot halo is difficult to reconcile with the relatively narrow line width observed for the NeVIII absorption feature,  $b_{\text{NeVIII}} = 23 \pm 15 \text{ km s}^{-1}$ , which corresponds to a velocity dispersion of  $\sigma_v = 16 \text{ km s}^{-1}$ . This is smaller than the velocity dispersion expected from virial motion along the line of sight  $\langle \sigma_v \rangle \equiv v_{\text{vir}} / \sqrt{3} \approx 62 \text{ km/s}$  and  $42 \text{ km/s}$  in a uniformly distributed hot gas in a halo of  $M_h = 10^{11.5} h^{-1} M_{\odot}$  (B) and  $M_h = 10^{11} h^{-1} M_{\odot}$  (A), respectively. We therefore do not favor this scenario.

#### 4.3. Photo-ionized Halo Gas

Next, we consider the possibility that the NeVIII absorber originates in individual clouds that are photo-ionized by both the intergalactic radiation field and local ionizing sources. Recall that Savage et al. (2005) ruled out a photo-ionization origin of the NeVIII absorber because the implied path length given a mean intergalactic radiation field of  $J_{912} = 1.9 \times 10^{-23} \text{ ergs cm}^{-2} \text{ s}^{-1} \text{ Hz}^{-1} \text{ sr}^{-1}$  at  $z = 0.2$  was  $\sim 11 \text{ Mpc}$ , too large for the observed line widths. The required path length would be smaller if the absorber was exposed to a more intense radiation source such as a local AGN. Here we examine whether our galaxy data can rule out the presence of such an ionizing source for producing the NeVIII absorber.

To reduce the path length to a more reasonable size ( $\sim 100$  kpc) requires the intensity of the radiation to increase by approximately two orders of magnitude (up to  $\sim 2 \times 10^{-21} \text{ erg cm}^{-2} \text{ s}^{-1} \text{ Hz}^{-1} \text{ sr}^{-1}$ ). Assuming a flat  $f_{\nu}$  spectrum (appropriate for an AGN) and adopting a  $r^{-2}$  scaling law, we estimate that an AGN would need to be as luminous as  $M_R - 5 \log h = -17.5$  at 38 kpc (A) or  $M_R - 5 \log h = -19.8$  at 110 kpc (B) away in order to provide sufficient ionizing photons for producing the NeVIII absorber<sup>2</sup>. The expected AGN luminosity already exceeds the observed luminosity for either of the two galaxies. In addition, there is a lack of AGN spectral features in the optical spectra. We therefore conclude that galaxies A and B are unlikely to host a powerful enough AGN to produce the observed absorption with photo-ionization.

While photo-ionization cannot explain the observed NeVIII absorber, we note that additional ionizing photons from galaxies A and B may contribute substantially to the ionizations of  $\text{H}^0$ ,  $\text{C}^{2+}$ ,  $\text{O}^{2+}$ , and  $\text{Si}^{2+}$ . Adopting a flat UV spectrum for star-forming galaxies and assuming 2% escape fraction of ionizing photons, we estimate the absorbing cloud is illuminated by additional ionizing radiation intensities of  $J_{912} = 1.3 \times 10^{-22}$  and  $0.3 \times 10^{-22} \text{ ergs cm}^{-2} \text{ s}^{-1} \text{ Hz}^{-1} \text{ sr}^{-1}$  from galaxies A and B, respectively. The inferred gas density and size of the H I absorbing cloud are then  $n_{\text{H}} = 2.2 \times 10^{-4} \text{ cm}^{-3}$  and  $\approx 4.7$

<sup>2</sup> Here we adopt  $H_0 = 70 \text{ km s}^{-1} \text{ Mpc}^{-1}$  for evaluating the distances between the absorber and galaxies A and B.




FIG. 1.— *Left*: Optical spectra of the three galaxies found within projected distances  $\rho < 200 h^{-1}$  kpc of the NeVIII absorber at  $z = 0.20701$  toward HE0226–4110. The spectra of galaxy A were obtained with LDSS3, while the spectra of galaxies B and C were obtained with IMACS (see Chen & Mulchaey 2009). The system response function of each slit setup has been removed based on a crude flux calibration using the spectra of alignment stars. *Right*: R-band image of the field around HE0226–4110. The seeing was  $\sim 0.65''$ .

kpc. By assuming a spherical cloud, we obtain a mass of  $M_{cl} \approx 6 \times 10^5 M_{\odot}$ . However, the mass could be larger with other assumed geometries.

#### 4.4. Outflows

Next, we consider the possibility that the absorption arises in a starburst driven wind from the dwarf galaxy (A). Based on the optical spectrum presented in Figure 1, we find that galaxy A is forming stars at a rate ( $\sim 0.1 M_{\odot} \text{ yr}^{-1}$ ) that is typical of dwarf galaxies at low redshift. With a rest-frame  $H\alpha$  equivalent width of  $EW \approx 13 \text{ \AA}$ , however, the galaxy would not be considered a starburst by most definitions adopted in the literature (Lee et al. 2009). Combining the estimated total stellar mass for galaxy A of  $M_* \sim 8 \times 10^8 h^{-2} M_{\odot}$  and the observed star formation rate, we derive a characteristic star formation time scale of  $\tau \approx 8 \text{ Gyr}$ . This is much longer than the characteristic time scale for typical starburst galaxies (Brinchmann et al. 2004). We also note that the major axis of the galaxy is oriented along the direction to the QSO. As an outflowing wind is expected to be predominantly along the minor axis, the bulk of the wind is unlikely to pass in front of the QSO

sightline for the observed geometry. Given the modest star formation rate in galaxy A and the disk alignment, a starburst driven wind would seem incapable of producing the observed absorption  $26.5 h^{-1}$  kpc from the disk of the galaxy.

#### 4.5. Conduction Front

Finally, we consider the idea that the NeVIII absorption is produced at the turbulent boundary layers of cooler clouds moving at high speed through a hot medium. This scenario has been proposed to explain the OVI absorption associated with high velocity clouds and the Magellanic Stream in the Milky Way (Sembach et al. 2003). The redshift difference between galaxy B and the NeVIII absorber leads to a relative velocity of nearly  $200 \text{ km s}^{-1}$ , roughly twice the sound speed for an ionized warm-hot medium of  $T \sim 5 \times 10^5 \text{ K}$ . Although the clouds are expected to suffer from the Kelvin-Helmholtz instability, the inferred gas density and cloud size discussed in § 4.3 suggest that the cloud may be massive enough to counterbalance such instabilities (Maller & Bullock 2004).

A natural consequence of this scenario is that both low and high ionization species would be expected with a moderate

velocity offset between the two components ( $\sim 20 \text{ km s}^{-1}$ ), similar to what is observed in the absorber at  $z = 0.20701$ . Therefore, the conductive front idea is both consistent with the observed galactic environment and the multi-phase nature of the gas.

## 5. CONCLUSIONS

Combining the results of our galaxy survey with the absorption line analysis of Savage et al. (2005), we have considered many possible origins for the  $z = 0.20701$  absorber toward HE0226–4110. We find that the absorber is best explained by a cool cloud moving through a hot medium. Although our study suggests that the NeVIII absorption may not be directly probing the WHIM in this system, it does imply the existence of hot gas at least  $76.5 h^{-1} \text{ kpc}$  from a  $0.25 L_*$  galaxy (B). The existence of extended hot gas halos around disk galaxies is a key ingredient in most semi-analytic models of disk galaxy formation (White & Frenk 1991; Cole et al. 2000; Bower et

al. 2006). However, there has been very little direct observational evidence for such halos in quiescent galaxies. Perhaps the best evidence for an extended hot corona comes from the Milky Way, where the presence of hot gas has been inferred from many different observations (Sembach et al. 2003; Fang et al. 2006; Bregman & Lloyd-Davies 2007). Our observations provide evidence for a similar halo around galaxy B. Given the large extent of the halo, a significant fraction of the baryons in the system are likely contained in hot gas.

We acknowledge valuable comments from the referee and useful discussions with Nick Gnedin, Juna Kollmeier, Janice Lee, Nicolas Lehner, Jesper Rasmussen, Josh Simon, Chris Thom and Rik Williams. H.-W.C. acknowledges partial support from NASA grant NNG06GC36G and NSF grant AST-0607510.

## REFERENCES

- Blanton, M. R., et al. 2003, *ApJ*, 592, 819  
 Bower, R. G., Benson, A. J., Malbon, R., Helly, J. C., Frenk, C. S., Baugh, C. M., Cole, S., & Lacey, C. G. 2006, *MNRAS*, 370, 645  
 Bregman, J. N. 2007, *ARA&A*, 45, 221  
 Bregman, J. N., & Lloyd-Davies, E. J. 2007, *ApJ*, 669, 990  
 Brinchmann, J., Charlot, S., White, S. D. M., Tremonti, C., Kauffmann, G., Heckman, T., & Brinkmann, J. 2004, *MNRAS*, 351, 1151  
 Cen, R., & Ostriker, J. P. 1999, *ApJ*, 514, 1  
 Chen, H.-W., & Mulchaey, J. S. 2009, *ApJ*, submitted  
 Cole, S., Lacey, C. G., Baugh, C. M., & Frenk, C. S. 2000, *MNRAS*, 319, 168  
 Danforth, C. W., & Shull, J. M. 2005, *ApJ*, 624, 555  
 Davé, R., et al. 2001, *ApJ*, 552, 473  
 Fabbiano, G., Kim, D.-W., & Trinchieri, G. 1992, *ApJS*, 80, 531  
 Fang, T., McKee, C. F., Canizares, C. R., & Wolfire, M. 2006, *ApJ*, 644, 174  
 Fukugita, M., Hogan, C. J., & Peebles, P. J. E. 1998, *ApJ*, 503, 518  
 Gnat, O., & Sternberg, A. 2007, *ApJS*, 168, 213  
 Haardt, F., & Madau, P. 1996, *ApJ*, 461, 20  
 Lee, J. C., Kennicutt, R. C., José G. Funes, S. J., Sakai, S., & Akiyama, S. 2009, *ApJ*, 692, 1305  
 Lehner, N., Savage, B. D., Wakker, B. P., Sembach, K. R., & Tripp, T. M. 2006, *ApJS*, 164, 1  
 Maller, A. H., & Bullock, J. S. 2004, *MNRAS*, 355, 694  
 Mulchaey, J. S., Mushotzky, R. F., Burstein, D., & Davis, D. S. 1996, *ApJ*, 456, L5  
 Persic, M., & Salucci, P. 1992, *MNRAS*, 258, 14P  
 Rasmussen, J., Sommer-Larsen, J., Pedersen, K., Toft, S., Benson, A., Bower, R. G., & Grove, L. F. 2009, arXiv:0903.0665  
 Savage, B. D., Sembach, K. R., Tripp, T. M., & Richter, P. 2002, *ApJ*, 564, 631  
 Savage, B. D., Lehner, N., Wakker, B. P., Sembach, K. R., & Tripp, T. M. 2005, *ApJ*, 626, 776  
 Sembach, K. R., et al. 2003, *ApJS*, 146, 165  
 Stocke, J. T., Penton, S. V., Danforth, C. W., Shull, J. M., Tumlinson, J., & McLin, K. M. 2006, *ApJ*, 641, 217  
 Thom, C., & Chen, H.-W. 2008a, *ApJ*, 683, 22  
 Thom, C., & Chen, H.-W. 2008b, *ApJS*, 179, 37  
 Tinker, J. L., & Conroy, C. 2009, *ApJ*, 691, 633  
 Tinker, J. L., Norberg, P., Weinberg, D. H., & Warren, M. S. 2007, *ApJ*, 659, 877  
 Tripp, T. M., Savage, B. D., & Jenkins, E. B. 2000, *ApJ*, 534, L1  
 Tripp, T. M., Sembach, K. R., Bowen, D. V., Savage, B. D., Jenkins, E. B., Lehner, N., & Richter, P. 2008, *ApJS*, 177, 39  
 Verner, D. A., Tytler, D., & Barthel, P. D. 1994, *ApJ*, 430, 186  
 Wakker, B. P., & Savage, B. D. 2009, arXiv:0903.2259  
 White, S. D. M., & Frenk, C. S. 1991, *ApJ*, 379, 52  
 Zehavi, I., et al. 2005, *ApJ*, 630, 1